

UCRL-JC-122218 Abs

**Penetration of Anthropogenic ^{12}C , ^{13}C , and ^{14}C into the ocean:
GCM simulations with implications for ocean uptake of bomb
radiocarbon**

Caldeira, Ken, Philip B. Duffy, & Greg H. Rau (Global Climate
Research Division, Lawrence Livermore National Laboratory, 7000
East Ave., Livermore, CA 94550 USA; 1-510-423-4191; internet:
kenc@llnl.gov)

We have developed a carbon cycle model that computes distributions of total inorganic ^{12}C , ^{13}C and ^{14}C , phosphate, alkalinity, and dissolved organic matter in the world ocean. This model is based on the work of Maier-Reimer, Najjar, Archer, and others. This biogeochemical model is coupled to a version of the GFDL ocean general circulation model that incorporates (1) a dynamic sea-ice model and (2) the Gent-McWilliams parameterization for the impact of isopycnal eddies on advection and diffusion in the ocean.

We will present the result of three pre-industrial to present model runs with different surface carbon and carbon isotope boundary conditions: (1) with observed atmospheric forcing; (2) with observed forcing less the inferred atmospheric bomb radiocarbon; and (3) with the pre-industrial atmosphere plus the inferred bomb radiocarbon.

We have two major conclusions. First, forcing ocean models with observed atmospheric $\Delta^{14}\text{C}$ values introduces significant errors into estimates of ocean bomb- ^{14}C uptake. Contrary to a common assumption, the number of atoms of bomb ^{14}C in the atmosphere is not directly proportional to atmospheric $\Delta^{14}\text{C}$, because of increasing concentrations of ^{12}C due to fossil fuel burning. Neglect of this fact results in underestimating the ocean bomb ^{14}C inventory after GEOSECS.

Second, neglect of increasing surface ocean ^{12}C concentrations has introduced a significant bias into the published estimates of ocean bomb ^{14}C concentrations based on the GEOSECS and other observations. These estimated bomb ^{14}C concentrations were made by subtracting $\Delta^{14}\text{C}$ values corresponding to estimated natural ^{14}C concentrations from observed $\Delta^{14}\text{C}$ values. Absent other anthropogenic influences (such as the Suess effect) on observed ocean $\Delta^{14}\text{C}$ values, this procedure would be correct. However, increasing ocean ^{12}C concentrations due to fossil fuel burning significantly affect observed ocean $\Delta^{14}\text{C}$ values. Neglect of this effect results in published inferred concentrations of ocean bomb ^{14}C being significantly underestimated.

This work was performed under the auspices of the U.S. Department of Energy by the Lawrence Livermore National Laboratory under Contract No. W-7405-Eng-48.

1. 1996 Ocean Sciences Meeting
2. 07737891
3. (a) Ken Caldeira, L-256, Lawrence
Livermore National Lab, P.O. Box
808, Livermore, Ca. 94550

(b) tel: (510) 423-4191
(c) fax: (510) 422-6388
4. 0
5. (a) None
(b) 1635, 4860, 4806
(c) N/A
6. 0
7. 0%
8. \$60 check enclosed
9. C
10. N/A
11. No